

Vortex state structure of a Bose condensate in an asymmetric trap

Anatoly A. Svidzinsky^{a,1}, Alexander L. Fetter^a

^a*Department of Physics, Stanford University, Stanford, CA 94305-4060*

Abstract

Based on an analytic solution of the Gross-Pitaevskii equation in the large-condensate (Thomas-Fermi) limit we determine the structure of a stationary vortex in a Bose-Einstein condensate in a nonaxisymmetric rotating trap. The condensate velocity field has cylindrical symmetry only near the vortex core and becomes intrinsically anisotropic near the condensate boundaries. Rotating the anisotropic trap induces an additional irrotational velocity field even for a vortex-free condensate.

Keywords: Bose condensate; vortices; nonaxisymmetric trap

The experimental achievement of Bose-Einstein condensation in confined alkali-atom gases [1–3] has stimulated great interest in the generation and observation of vortices in such systems [4–6]. Rotating a totally anisotropic harmonic trap at an angular frequency Ω can, in principle, generate vortices; they are energetically stable for $\Omega > \Omega_c$ [7–10].

This work determines the condensate velocity field for a stationary vortex when the completely anisotropic harmonic trap potential $V_{\text{tr}} = \frac{1}{2}M(\omega_x^2 x^2 + \omega_y^2 y^2 + \omega_z^2 z^2)$ rotates about a principal axis. We consider a zero-temperature condensate in the Thomas-Fermi (TF) limit, when the vortex core radius $\xi \sim d^2/R$ is small compared to the mean oscillator length d and the mean dimension R of the condensate [here, $d = (d_x d_y d_z)^{1/3}$

with $d_i = \sqrt{\hbar/M\omega_i}$ and trap frequencies ω_i ($i = x, y, z$)].

We assume that the trap rotates with an angular velocity Ω around the z axis. At zero temperature in the rotating frame, the equilibrium condensate wave function Ψ satisfies the Gross-Pitaevskii equation:

$$\left(-\frac{\hbar^2}{2M}\nabla^2 + V_{\text{tr}} + g|\Psi|^2 - \mu + i\hbar\Omega\partial_\phi\right)\Psi = 0, \quad (1)$$

where $g = 4\pi\hbar^2 a/M > 0$ is the effective interparticle interaction strength, μ is the chemical potential in the rotating frame, and ϕ is the azimuthal angle in cylindrical polar coordinates (r, ϕ, z) . Consider a q -fold vortex, oriented along the z axis, with a condensate wave function $\Psi = |\Psi|e^{iS}$. Its phase must have the form $S = q\phi + S_1$, where S_1 is an odd 2π -periodic function of ϕ .

Multiplying the imaginary part of (1) by $|\Psi|$ yields the equation of current continuity in the rotating frame:

¹ Corresponding author.
asvid@leland.Stanford.EDU

E-mail:

$$\text{div}(|\Psi|^2 \nabla S) - \frac{M}{\hbar} \Omega \partial_\phi |\Psi|^2 = 0. \quad (2)$$

The second term here arises from the difference between the condensate velocity field $\mathbf{v}^l = \frac{\hbar}{M} \nabla S$ in the laboratory frame and $\mathbf{v}^r = \frac{\hbar}{M} \nabla S - \Omega \times \mathbf{r}$ in the rotating frame. In the TF limit in the absence of a vortex, $g|\Psi_{TF}|^2 = \mu - V_{\text{tr}}$, and for distances far from the vortex core ($r \gg |q|\xi$) we may substitute $|\Psi_{TF}|$ into (2) instead of $|\Psi|$, getting

$$\frac{2g}{M} |\Psi_{TF}|^2 \nabla^2 S - [\omega_x^2 + \omega_y^2 + (\omega_x^2 - \omega_y^2) \cos(2\phi)] r \partial_r S + (\omega_x^2 - \omega_y^2) \sin(2\phi) \left(\partial_\phi S - \frac{M\Omega}{\hbar} r^2 \right) - 2\omega_z^2 z \partial_z S = 0.$$

Let us seek a solution of this equation in the form:

$$S = S_0 + q\phi - \beta\Omega r^2 \sin(2\phi), \quad (3)$$

where $\beta = (M/2\hbar) (\omega_x^2 - \omega_y^2) / (\omega_x^2 + \omega_y^2)$. The resulting equation for S_0 does not contain Ω . Further, let us consider distances $x, y, z \ll R_x, R_y, R_z$, where R_i are the TF dimensions of the condensate ($M\omega_i^2 R_i^2 = 2\mu$, $i = x, y, z$). At these intermediate distances we can omit terms of higher order in the parameter $(x_i/R_i)^2$. As a result, we find the following inhomogeneous equation:

$$\nabla^2 S_0 + \left(\frac{1}{R_x^2} - \frac{1}{R_y^2} \right) q \sin(2\phi) = 0. \quad (4)$$

This equation has an explicit periodic solution

$$S_0 = -\alpha q r^2 \ln\left(\frac{r}{A}\right) \sin(2\phi), \quad (5)$$

where $\alpha = \frac{1}{4} (R_x^{-2} - R_y^{-2})$ and A is a constant of integration that must be chosen to minimize the total energy. We estimate $A \sim R_\perp$, where $R_\perp^2 = 2R_x^2 R_y^2 / (R_x^2 + R_y^2)$. Thus, the phase S near the center of a rotating trap in the presence of a vortex is

$$S \approx q\phi - \left[\alpha q \ln\left(\frac{r}{R_\perp}\right) + \beta\Omega \right] r^2 \sin(2\phi). \quad (6)$$

Near the vortex core, the condensate wave function and condensate velocity possess cylindrical symmetry, while far from the vortex core the condensate velocity adjusts to the anisotropy of the

trap and becomes asymmetric. The condensate velocity in the laboratory frame becomes

$$\frac{M\mathbf{v}^l}{\hbar} \approx \left\{ \frac{q}{r} - 2 \left[\alpha q \ln\left(\frac{r}{R_\perp}\right) + \beta\Omega \right] r \cos(2\phi) \right\} \mathbf{e}_\phi - 2 \left[\alpha q \ln\left(\frac{r}{R_\perp}\right) + \beta\Omega \right] r \sin(2\phi) \mathbf{e}_r. \quad (7)$$

The energy barrier to vortex formation [9] means that the condensate can be vortex free even for $\Omega > \Omega_c$, in which case the trap rotation induces an irrotational velocity field $\mathbf{v}^l \propto \Omega \nabla(xy)$ [11]. To logarithmic accuracy, the additional phase S_0 does not affect the critical angular velocity Ω_c and metastable angular velocity $\Omega_m = \frac{3}{5}\Omega_c$ found previously [9].

In conclusion, an analytic solution of Gross-Pitaevskii equation in the rotating frame gives the condensate velocity field for a vortex state in an anisotropic rotating trap. Even for a vortex-free condensate, the rotation induces superfluid motion.

This work was supported in part by the National Science Foundation, Grant No. DMR 94-21888, and by Stanford University (A.A.S.).

References

- [1] M. H. Anderson *et al.*, Science **269**, 198 (1995).
- [2] C. C. Bradley *et al.*, Phys. Rev. Lett. **75**, 1687 (1995).
- [3] K. B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995).
- [4] B. Jackson, J. F. McCann, and C. S. Adams, Phys. Rev. Lett. **80**, 3903 (1998).
- [5] R. Dum *et al.*, Phys. Rev. Lett. **80**, 2972 (1998).
- [6] K. P. Marzlin, W. Zhang, and E. M. Wright, Phys. Rev. Lett. **79**, 4728 (1997).
- [7] G. Baym and C. J. Pethick, Phys. Rev. Lett. **76**, 6 (1996).
- [8] F. Dalfovo and S. Stringari, Phys. Rev. A **53**, 2477 (1996).
- [9] A. A. Svidzinsky and A. L. Fetter, e-print: cond-mat/9811348.
- [10] D. L. Feder, C. W. Clark, and B. I. Schneider, e-print: cond-mat/9904269.
- [11] A. L. Fetter, J. Low Temp. Phys. **16**, 533 (1974).